

STOCHASTIC INVERTIBLE MAPPINGS BETWEEN POWER LAW AND GAUSSIAN PROBABILITY DISTRIBUTIONS

C. VIGNAT¹ AND A. PLASTINO²

¹ L.I.S. GRENOBLE, FRANCE AND E.E.C.S., UNIVERSITY OF MICHIGAN, USA
E-MAIL: VIGNAT@UNIV-MLV.FR

² LA PLATA NATIONAL UNIVERSITY & NATIONAL RESEARCH COUNCIL (CONICET)
C. C. 727 - 1900 LA PLATA - ARGENTINA
E-MAIL: PLASTINO@FISICA.UNLP.EDU.AR

ABSTRACT. We construct “stochastic mappings” between power law probability distributions (PD’s) and Gaussian ones. To a given vector N , Gaussian distributed (respectively Z , exponentially distributed), one can associate a vector X , “power law distributed”, by multiplying X by a random scalar variable a , $N = aX$. This mapping is “invertible”: one can go via multiplication by another random variable b from X to N (resp. from X to Z), i.e., $X = bN$ (resp. $X = bZ$). Note that all the above equalities mean “is distributed as”. As an application of this stochastic mapping we revisit the so-called “zero-th law of thermodynamics problem” that bedevils the practitioners of nonextensive thermostatics.

Keywords: Superstatistics, stochastic mappings, Tsallis entropies

PACS: 05.40.-a, 05.20.Gg

1. INTRODUCTION

Statistical Mechanics’ most notorious and renowned probability distribution (PD) is that deduced by Gibbs for the canonical ensemble [1, 2], usually referred to as the Boltzmann-Gibbs (BG) equilibrium distribution

$$(1) \quad p_G(i) = \frac{\exp(-\beta E_i)}{Z_{BG}},$$

with E_i the energy of the microstate labeled by i , $\beta = 1/k_B T$ the inverse temperature (T), k_B Boltzmann’s constant, and Z_{BG} the partition function. The exponential term $F_{BG} = \exp(-\beta E)$ is, of course, called the BG factor. Recently Beck and Cohen [3] have advanced a generalization (called “superstatistics”) of this BG factor, assuming that the inverse temperature β is a stochastic variable. They effect a multiplicative convolution to obtain a generalized statistical factor F_{GS} in the fashion

$$(2) \quad F_{GS} = \int_0^\infty \frac{d\beta}{\beta} f(\beta) \exp(-\beta E) \equiv f \circ F_{BG},$$

where $f(\beta)$ satisfies

$$(3) \quad \int_{-\infty}^\infty d\beta f(\beta) = 1.$$

Note that the $\circ$ -sign is used to denote the multiplicative convolution between two PDs: multiplicative convolution of two PDs f_X and f_Y is the PD of the product of the two corresponding random variables X and Y .

As stated above, β is the inverse temperature, but the integration variable may also be any convenient intensive parameter. Superstatistics would arise as a “statistics of statistics” that takes into account fluctuations of such intensive parameters. Beck and Cohen also show that, if $f(\beta)$ is a χ^2 distribution, nonextensive thermostatics (NEXT) is obtained, which is of interest because NEXT, or Tsallis’ thermostatics, is today a very active field, perhaps a new paradigm for statistical mechanics, with applications to several scientific disciplines [4, 5, 6]. In working in a NEXT framework one has to deal with power-law distributions, which are certainly ubiquitous in physics (critical phenomena are just a conspicuous example [7]). Indeed, it is well known that power-law distributions arise quite naturally in maximizing Tsallis’ information measure (q is a real parameter called the “nonextensivity index”)

$$(4) \quad H_q(f) = \frac{1}{1-q} \left(1 - \int_{-\infty}^{+\infty} f(x)^q dx \right),$$

subject to appropriate constraints. In the case of the canonical distribution there is only one constraint, the energy E , i.e., $\langle E \rangle = K$ (K a constant) and the equilibrium canonical distribution writes $f(x) = (1/Z_q)[1 - (1-q)\beta_q E]^{\frac{1}{q-1}}$, with β_q and Z_q standing for the NEXT counterparts of β and Z_{BG} above. It is a classical result that as $q \rightarrow 1$, Tsallis entropy reduces to Shannon entropy

$$(5) \quad H_1(f) = - \int_{-\infty}^{+\infty} f(x) \log f(x).$$

Tsallis’ PDs are encountered in analyzing a rather large variety of physical systems [8, 9, 10, 11, 12], which encourages people to continue investigating the nonextensive formalism along multiple viewpoints and a multitude of paths.

In such a spirit, and further pursuing along the road first travelled by Beck and Cohen [3], we show here that, for a fixed temperature T (respectively, any adequate intensive parameter τ), *there exists a mapping between power law PD’s on the one hand, and exponential or Gaussian PD’s on the other one*. Using this mapping one can transform a Tsallis PD into a Gaussian (resp., exponential) PD and vice-versa via multiplicative convolution with a chi random variable of variance unity.

This mapping nitidly reveals, for the first time as far as we know, the (perhaps surprising) intimate relation that exists between the conventional Shannon-Gibbs-Boltzmann statistics and NEXT. As a first application we will tackle the vexing (for NEXT) problem that revolves around thermodynamics zero-th law that was first revealed by Raggio and Guerberoff [13] (see also [14, 15, 16, 17, 18], a by no means exhaustive list, and references therein).

2. MATHEMATICAL FRAMEWORK

2.1. Case $q > 1$. Let us consider a random k -variate vector $X = (x_1, \dots, x_k)^T$ following a Tsallis distribution with parameter $q > 1$ (or $q < 0$). Its distribution writes

$$(6) \quad f_X(X) \propto (1 - X^T X)_+^{\frac{n-k}{2}-1}$$

with $n \geq k$ and with the notation $x_+ = \max(0, x)$. As discussed in [19], f_X can be obtained as the k -variate marginal of a uniform distribution on the sphere in $\mathbb{R}^n$. Additionally, a stochastic representation for X can be given in terms of Gaussians, which is essential for our present purposes, i.e.,

$$(7) \quad X = \frac{\tilde{N}}{r}$$

where $N = [N_1, \dots, N_n]^T$ is a Gaussian n -variate vector, $\tilde{N}$ is the k -variate vector composed of the first k components of N and $r = \sqrt{\sum_{i=1}^n N_i^2}$ is a chi random variable with n degrees of freedom. We note that the first k components of N are shared by the numerator and the denominator.

Theorem. *If a_n is i) a chi random variable with n degrees of freedom and ii) independent of a vector X distributed as in (6), then the product*

$$(8) \quad a_n X \sim Z$$

where Z is Gaussian with identity covariance and symbol $\sim$ denotes equality in distribution.

Proof. See the Annex. □

Thus a Tsallis system can be "Gaussianized" simply by multiplying each of its components by an independent, scalar chi random variable with n degrees of freedom. For statistical mechanics purposes, the variance is a very important quantity because it is proportional to the temperature [2]. As defined by (6), the random vector X has covariance matrix $R_X = \frac{1}{n}I_k$, with I_k the $(k \times k)$ -unit matrix. Thus, the random vector Z , as defined by (8) has covariance matrix (we use the notation $EX \equiv \langle X \rangle$)

$$(9) \quad EZZ^T = Ea_n^2 EXX^T = n \times \frac{1}{n}I_k = I_k.$$

2.2. Case $\frac{k}{k+1} < q < 1$. Let us first underline a useful connexion between the instances $q < 1$ and $q > 1$:

Lemma. *if X is distributed according to (6) then*

$$(10) \quad Y = \frac{X}{\sqrt{1 - X^T X}}$$

is distributed as follows:

$$(11) \quad f_Y(Y) \propto (1 + Y^T Y)^{-\frac{m+k}{2}}$$

with $m = n - k$, and thus is a k -variate Tsallis random vector with parameter $\frac{k}{k+2} < q < 1$. A stochastic representation of Y writes

$$(12) \quad Y = \frac{N}{\chi_m},$$

where N is a k -variate Gaussian vector and χ_m is a chi random variable independent on N with $m = n - k$ degrees of freedom. Note moreover that the covariance matrix of Y writes

$$(13) \quad EYY^T = ENN^T E \frac{1}{\chi_m^2} = \frac{1}{m-2}I_k.$$

Now we can state our main theorem:

Theorem. *if Y is distributed according to (11) and if a_n is a chi distributed random variable with n degrees of freedom independent on Y , then random variable*

$$(14) \quad Z = \frac{a_n}{\sqrt{1 + Y^T Y}} Y$$

is Gaussian with unit covariance. Moreover, random variable

$$(15) \quad b_m = \frac{a_n}{\sqrt{1 + Y^T Y}},$$

is chi distributed with m degrees of freedom. Note that b_m is dependent on Y .

Proof. See the Annex. □

2.3. Extension of the above theorems. The precedent considerations can be extended from the Gaussian case to the one of a purely exponential factor. A given Tsallis-distributed vector X gets transformed, via multiplicative convolution with a chi random variable, into a vector that is distributed according to an exponential PD. In other words, from a Gaussian vector G , one can create a Tsallis distributed vector X by multiplying by a random scalar variable. This mapping is invertible: a Gaussian distributed vector can be obtained by multiplying a Tsallis distributed one by another properly chosen random variable.

2.4. Application: the merging of two independent systems. Suppose we have two independent systems (in the sense that their mutual interaction is negligible) whose states are described by two Tsallis random vectors, independent as well, X and Y with, say, $q > 1$. If we consider the system $Z^T = [X^T, Y^T]$ we immediately realize that it is not Tsallis distributed since its distribution writes

$$(16) \quad f_Z(X, Y) = (1 - X^T X)^{\frac{n-k}{2}-1} (1 - Y^T Y)^{\frac{n-k}{2}-1},$$

and, consequently, can not be expressed as a function of $X^T X + Y^T Y$ (except in the Gaussian case $q = 1$) [13, 14, 15, 16, 17].

The problem can, however, be circumvented as follows: if we now "pre-multiply" system X (resp. Y) with the same chi-distributed random variable a_n as defined in theorem 2.1, then the system

$$(17) \quad Z = a_n [X^T, Y^T]^T$$

is the merging of two Gaussian systems. Moreover, the covariance of Z writes

$$(18) \quad E Z Z^T = I_{2k},$$

so that $a_n X$ and $a_n Y$ are both uncorrelated and Gaussian, and thus independent. Finally, considering a new chi distributed random variable b_m with m degrees of freedom and independent of Z , the Tsallis distributed vector

$$(19) \quad X_{Tsallis} = \frac{Z}{b_m} = \frac{a_n}{b_m} [X^T, Y^T]^T,$$

- and NOT (16) - turns out to be the "true" representative of the "merging" of two nonextensive systems. Such a vector is not characterized by a fixed temperature T (or, more generally, by a fixed value of an appropriate intensive system's parameter τ), but by a superposition of temperatures "centered" at T (resp., superposition

of the intensive parameter centered at τ), exactly in the spirit of superstatistics. We can then speak of a “stochastic” zero-th law that would apply for nonextensive thermostatics: *given two independent systems A , B of equal temperature T , the pertinent temperature of the associated composite system $A + B$ at equilibrium fluctuates around T .*

3. PHYSICAL CONSIDERATIONS

For didactic reasons it is convenient at this point to briefly review the well-known connection between variance and temperature for a Gaussian PD before probing further into the zero-th law.

3.1. Quadratic classical Hamiltonians. We consider first the “normal modes” hamiltonian for N independent particles in three dimensions

$$(20) \quad H = \frac{1}{2} \sum_{m=1}^{3N} c_m^q q_m^2 + c_m^p p_m^2 \equiv \sum_{i=1}^{6N} c_i z_i^2 / 2,$$

and consider the location of the many-body system in its $6N$ –dimensional Γ –space. This is represented by a $6N$ –vector. Obviously, Γ –position has probabilities proportional to a function of βH . We consider two instances: conventional BG thermostatics and Tsallis’ one.

- For BG statistics, location is represented by a vector Z distributed according to the $(1/Z) \exp(-\beta H)$ PD. On account of (20) then, Z is Gaussian-distributed.
- For Tsallis’ statistics, locations is represented by a vector X that is Tsallis-distributed.

If $Z = (z_1, \dots, z_{6N})^T$ is BG-distributed, then its covariance matrix EZZ^T is a diagonal matrix. Because of the equipartition theorem [1], the expectation value of each member of the diagonal equals $k_B T/2$. Thus, $EZZ^T = (k_B T/2)I_{6N}$, and the covariance matrix is proportional to the temperature.

In the case of Tsallis statistics, let the vectors in (16) be distributed according to

$$(21) \quad \begin{aligned} f_X(x) &\propto [1 - (1 - q)\beta \sum_{i=1}^k x_i^2] \\ f_Y(y) &\propto [1 - (1 - q)\beta \sum_{i=1}^k y_i^2], \end{aligned}$$

so that we are merging two system of inverse temperature $\beta = 1/k_B T$. The representative of the merged system is (19). Note that Z there is a Gaussian vector of temperature T , and $X_{Tsallis}$ a Tsallis vector that, although not having a definite temperature, possesses a temperature distribution “peaked” at T . The zero-th law of thermodynamics for Tsallis distributed vectors entails that if we join two nonextensive systems of temperature T and wait until equilibrium is re-established, we obtain a total system whose temperature “fluctuates” (à la Beck-Cohen) around this temperature T .

3.2. “Normal modes” or diagonalizable Hamiltonians. Our approach can be extended to more general types of Hamiltonians of the type

$$(22) \quad H = \sum_{i=1}^{6N} c_i z_i^{p_i}, \quad p_i \text{ an integer power},$$

provided the mixing variable a_n is modified as follows.

Theorem. *If A is a diagonal matrix whose element $A_{i,i} = a^{\frac{1}{p_i}}$ where a is i) a chi random variable with $n - k + 2 \sum_{i=1}^k \frac{1}{p_i}$ degrees of freedom and ii) independent of component X_i distributed as in (6), then the product*

$$(23) \quad AX \sim Z$$

where Z is distributed as

$$(24) \quad f_Z(Z) \propto \exp \left(- \sum_{i=1}^k c_i z_i^{p_i} \right).$$

Proof. See the Annex. □

Following now an identical line of reasoning as that of Subsection 2.4, we can generalize the stochastic zero-th law obtained there to all classical systems described by the above type of Hamiltonian.

4. CONCLUSIONS

We have here generalized the concept of superstatistics devised by Beck and Cohen [3]. Its typical “convolution path” that involves fluctuations of intensive parameters allows one to express generalized factors B_{GF} in terms of a superposition of exponential PDs. We show the the path can be “inverted”, at least with reference to Tsallis PDs, in the sense that a normal (or an exponential) distribution can also be cast as a superposition of Tsallis PD. *Stochastic invertible mappings between power law distributions and Gaussian (or exponential) ones have been introduced and discussed.*

As an application we have shown that the zero-th law of thermodynamics is obeyed by Tsallis PDs only in a stochastic sense. In nonextensive thermostatics, if two systems of temperature T are merged, the ensuing equilibrium total distribution exhibits fluctuation around that temperature.

5. ANNEX: PROOFS

5.1. Proof of theorem 2.1. From [25, 3.471.3], we have

$$(25) \quad \exp(-cx) = \frac{c^\alpha}{\Gamma(\alpha)} \int_0^{+\infty} \exp(-c/u) u^{-\alpha-1} (1-ux)_+^{\alpha-1} du$$

Now replace x by $\sum_{i=1}^k x_i^2$ and u by v^{-1} in (25) to obtain

$$\begin{aligned} \exp \left(-c \sum_{i=1}^k x_i^2 \right) &= \frac{c^\alpha}{\Gamma(\alpha)} \int_0^{+\infty} \exp(-cv) v^{\alpha-1} \left(1 - v^{-1} \sum_{i=1}^k x_i^2 \right)_+^{\alpha-1} dv \\ &= \frac{c^\alpha}{\Gamma(\alpha)} \int_0^{+\infty} \exp(-cv) v^{\alpha-1+\frac{k}{2}} \left(\frac{1}{\sqrt{v}} \right)^k \left(1 - \sum_{i=1}^k \frac{x_i^2}{v} \right)_+^{\alpha-1} dv. \end{aligned}$$

Now, as a classical Gamma integral,

$$\int_0^{+\infty} \exp(-cv) v^{\alpha-1+\frac{k}{2}} dv = \frac{\Gamma(\alpha + k/2)}{c^{\alpha+k/2}}$$

so that

$$f_v(v) = \frac{c^{\alpha+k/2}}{\Gamma(\alpha + k/2)} \exp(-cv) v^{\alpha-1+\frac{k}{2}}$$

is the distribution of a chi-squared random variable with $k + 2\alpha$ degrees of freedom and variance $\sigma^2 = 1/2c$. Thus

$$\begin{aligned} \left(\frac{1}{2\pi\sigma^2}\right)^{k/2} \exp\left(-\frac{\sum_{i=1}^k x_i^2}{2\sigma^2}\right) &= \frac{c^\alpha \Gamma(\alpha + k/2)}{c^{\alpha+k/2} (\sigma\sqrt{2\pi})^k \Gamma(\alpha)} \int_0^{+\infty} \sigma \chi_{2\alpha+k}(v) \left(\frac{1}{\sqrt{v}}\right)^k \left(1 - \sum_{i=1}^k \frac{x_i^2}{v}\right)_+^{\alpha-1} dv. \\ &= \int_0^{+\infty} \sigma \chi_{2\alpha+k}(v) S_{X\sqrt{v}}^{(\alpha)}(x_1, \dots, x_k) dv \end{aligned}$$

where $S_X^{(\alpha)}$ denotes a k -variate Tsallis distribution obtained as the marginal of the uniform distribution on the sphere S_n

$$S_X^{(\alpha)}(x) = \frac{\Gamma(\alpha + k/2)}{(2\pi\sigma^2 c)^{\frac{k}{2}} \Gamma(\alpha)} (1 - x^T x)^{\alpha-1}$$

where

$$\frac{n-k}{2} = \alpha \leftrightarrow n = 2\alpha + k.$$

5.2. Proof of theorem 2.2. We know from the lemma that

$$Y = \frac{X}{\sqrt{1 - X^T X}}$$

and we deduce that

$$X = \frac{Y}{\sqrt{1 + Y^T Y}}.$$

From theorem 2.1, we know that $a_n X$ is normal. Thus

$$\frac{a_n}{\sqrt{1 + Y^T Y}} Y$$

is normal. Moreover, the distribution of random variable $b_m = a_n/\sqrt{1 + Y^T Y}$ can be computed as follows

$$\begin{aligned} f_b(b) &\propto \int (1 + Y^T Y)^{n/2} b^{n-1} \exp\left(-\frac{b^2(1 + Y^T Y)}{2}\right) (1 + Y^T Y)^{-\frac{n+k}{2}} dY \\ &= b^{n-1} \exp\left(-\frac{b^2}{2}\right) \int \exp\left(-\frac{b^2 Y^T Y}{2}\right) dY \\ &\propto b^{n-1} \exp\left(-\frac{b^2}{2}\right) b^{-k} = b^{m-1} \exp\left(-\frac{b^2}{2}\right) \end{aligned}$$

so that b_m is a chi random variable with m degrees of freedom.

5.3. **Proof of theorem 3.2.** Starting from equality (25), we obtain

$$\begin{aligned} \exp\left(-c \sum_{i=1}^k x_i^{p_i}\right) &= \frac{c^\alpha}{\Gamma(\alpha)} \int_0^{+\infty} \exp(-cv) v^{\alpha-1} \left(1 - \sum_{i=1}^k \left(\frac{x_i}{v^{\frac{1}{p_i}}}\right)^{p_i}\right)_+^{\alpha-1} dv \\ &= \frac{c^\alpha}{\Gamma(\alpha)} \int_0^{+\infty} \exp(-cv) v^{\alpha-1+\frac{1}{p}} \left(\prod_{i=1}^k \frac{1}{v^{\frac{1}{p_i}}}\right) \left(1 - \sum_{i=1}^k \left(\frac{x_i}{v^{\frac{1}{p_i}}}\right)^{p_i}\right)_+^{\alpha-1} dv \end{aligned}$$

with $\frac{1}{p} = \sum_{i=1}^k \frac{1}{p_i}$. Following the same path as in proof of theorem 3.2, we deduce that vector with i -th component $a^{p_i} X_i$, where X_i is distributed as

$$(26) \quad f_{X_i}(x) = (1 - x^{p_i})_+^{\alpha-1},$$

is distributed as vector Z with

$$(27) \quad f_Z(Z) \propto \exp\left(-\sum_{i=1}^k c_i z_i^{p_i}\right)$$

REFERENCES

- [1] Reif F 1965 *Statistical and thermal physics* (NY: McGraw-Hill); Pathria R K 1993 *Statistical Mechanics* (Exeter: Pergamon Press).
- [2] Gibbs J W 1948 *Elementary principles in statistical mechanics in Collected Works* (New Haven: Yale University Press); R. B. Lindsay R B and Margenau H 1957 *Foundations of physics* (NY: Dover).
- [3] C. Beck and E. G. D. Cohen, *Physica A* **322** (2003) 267.
- [4] Gell-Mann M and Tsallis C, Eds. 2004 *Nonextensive Entropy: Interdisciplinary applications* (Oxford: Oxford University Press) and references therein; Plastino A R and Plastino A 1994 *Phys. Lett. A* **193** 140.
- [5] Kaniadakis G, Lissia M, and Rapisarda A., Eds., 2002 *Nonextensive statistical mechanics and physical applications*, *Physica A (Special)* **305**, and references therein.
- [6] A.R. Plastino, A. Plastino, *Phys. Lett. A* **193** (1994) 251.
- [7] N. Goldenfeld, *Lectures on phase transitions and the renormalization group* (Addison-Wesley, NY, 1992).
- [8] C. Tsallis, *Braz. J. Phys.* **29** (1999) 1.
- [9] C. Beck, *Phys. Lett. A* **287** (2001) 240.
- [10] C. Beck, G. S. Lewis, and H. L. Swinney, *Phys. Rev. E* **63** (2001) 035503.
- [11] I. Bediaga, E. M. F. Curado, and J. M. de Miranda, *Physica A* **286** (2000) 156.
- [12] C. Tsallis, J. C. Anjos, and E. P. Borges, *Phys. Lett. A* **310** (2003) 372.
- [13] G. R. Guerberoff and G. A. Raggio, *J. Math. Phys.* **37** (1996) 1776.
- [14] C. Tsallis, "Nonextensive statistical mechanics and thermodynamics. Historical background and present status" in *Nonextensive statistical mechanics and its applications*, Eds. S. Abe and Y. Okamoto ("Lecture Notes in Physics", Springer, Berlin, 2000).
- [15] S. Martinez, F. Pennini, and A. Plastino, *Physica A* **295** (2001) 224.
- [16] Q. A. Wang, L. Nivenen, A. Le Méhauté, and M. Perezil, *J. Phys. A* **35** (2002) 7003.
- [17] S. Abe, S. Martinez, F. Pennini, and A. Plastino, *Phys. Lett. A* **281** (2001) 126.
- [18] S. Abe, *Physica A* **269** (1999) 403.
- [19] C. Vignat, A. Plastino, cond-mat/0503337 *Title: Geometric origin of probabilistic distributions in statistical mechanics.*
- [20] Scalapino D J 1993 *Physics and probability. Essays in honor of Edwin T. Jaynes, edited by Grandy W T, Jr. and Milonni P W* (NY: Cambridge University Press), and references therein.
- [21] Jaynes E T 1957 *Phys. Rev.* **106** 620; **108** 171; 1987 *Papers on probability, statistics and statistical physics, edited by Rosenkrantz R. D.* (Dordrecht: Reidel); Katz A 1967 *Principles of Statistical Mechanics: The Information Theory Approach* (San Francisco: Freeman and Co.).

- [22] Poincaré H 1912 *Calcul des probabilités* (Paris: Gauthier-Villars).
- [23] McKean H P 1973 Ann. Probability **1** 197.
- [24] Sakurai J J 1985 *Modern quantum mechanics* (Menlo Park, Ca.: Benjamin).
- [25] Gradshteyn and Ryzhik's Table of Integrals, Series, and Products, Alan Jeffrey (ed.), Fifth edition (January 1994)